



## Effect of Variegated Forest Soil Amendments on the Germination and Early Growth of *Irvingia gabonensis* (O Rorke, Baill)

A. E. Egwunatum<sup>1\*</sup>, D. E. Dolor<sup>2</sup> and C. J. Ofobike<sup>1</sup>

<sup>1</sup>Department of Forestry and Wildlife, Nnamdi Azikiwe University, Awka, Anambra State, Nigeria.

<sup>2</sup>Department of Agronomy, Forestry and Wildlife, Delta State University, Abraka, Delta State, Nigeria.

### Authors' contributions

This work was carried out in collaboration among all authors. Author AEE designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors DED and CJO managed the analyses of the study. Author DED managed the literature searches. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/AJRAF/2020/v6i130098

#### Editor(s):

(1) Dr. Lucia Bortolini, University of Padova, Italy.

#### Reviewers:

(1) Anwar Masood, MJP Rohilkhand University, India.

(2) Prawal Pratap Singh Verma, CIMAP Research Centre, India.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/58259>

Original Research Article

Received 12 April 2020

Accepted 18 June 2020

Published 24 June 2020

### ABSTRACT

This study evaluated the early germination and growth variables of *Irvingia gabonensis* under organically primed and thermally amended soil media. Six media were prepared from Gmelina plantation topsoil by amendment with poultry waste (T2), river sand (T3) and combusted forest floor litters for 5 (T4), 10 (T5), 15 (T6) and 20 (T7) minutes respectively to contrast a control topsoil (T1). Soil media were analyzed for critical nutrient properties and engaged in the germination and early growth trial of *Irvingia* replicated three times, and arranged in a completely randomized design. Data collected were analyzed with ANOVA and significant means separated with the Duncan multiple range test. The results showed that Ca/Mg ratio was widest for T3, least CEC (13.2 meq/100 g soil) by T7 and slightly acidic pH (H<sub>2</sub>O) for T1 and T2 while T3, T4, T6 and T7 were alkaline. Germination at 6WAS was T4 (100%) > T1=T2=T5 (85.71%) > T3 (71.43%) > T6 (42.86%) > T7 (0%). The growth variables of seedling varied significantly (P<0.05) with T3 and T4 comparing favorably in collar diameter (4.50x10<sup>-1</sup>± 0.05 mm) and leaf area (35.08±4.85 mm<sup>2</sup>) although T3 recorded the highest stem height (117.79±0.42 mm). The study recommends the use of least

\*Corresponding author: E-mail: ae.egwunatum@unizik.edu.ng;

thermally modified media (T4) and primed topsoil-riversand (T3) for fast germination in view of conserving food reserve for the critical early growth period of *I. gabonensis* in pursuit of its domestication in nutrient degraded soils.

**Keywords:** Thermal soil amendment; organic priming; topsoil; cation exchange capacity (CEC); *Irvingia gabonensis*.

## 1. INTRODUCTION

The enrichment of forest soils for the growth of forest tree species is intrinsically receiving attention as way of reducing the long gestation periods of these species in view of domestication. This approach is significant as the original natural forest with its succession capability is currently under threat due to various anthropogenic activities especially the proximate drivers of deforestation [1,2,3]. Consequently, a good proportion of indigenous forest tree species have become threatened on the IUCN list as a result of deforestation and the inadequate management of declining forest soil resource alongside observed threats which potentially jeopardize attempts to effectively domesticate before disappearance in the wild. This is because the hitherto forest soils under luxuriant vegetation were often enriched with litters of twigs, leaves and other component bi-products of natural interaction within the forest to provide capacity for fallen seeds to germinate in the wild, and where this capacity is lost, the germination potential of seeds as well as growth of seedlings become critical as the food reserves contained in the cotyledons declines with prolonged, retarded early growth [4].

*Irvingia gabonensis* also known as dika nuts belong to the Irvingiaceae family. It is an evergreen tree species with sufficiently dense crown that can grow up to 15-40 m tall and girth not less than 100 cm [5] and commonly distributed on a hotspot belt that runs from Nigeria in West Africa through Central to South Africa. The tree produces yellowish varieties that are either edible fibrous fruit with turpentine flavor or bitter acrid taste that have various industrial, commercial and medicinal values [6,7]. The tough, heavy and tannin-immune wood content makes *Irvingia* less susceptible to bio deterioration and increases its susceptibility as choice tree species for exploitation as timber.

Currently, *Irvingia gabonensis* has been classified as "Near threatened" in the list of threatened species [8] due to logging operations

and poor natural regeneration which may not be unconnected with degraded soil nutrients owing to loss in vegetation vis-à-vis low litter falls. It is therefore critical to source for appropriate soil media through modification with available organic matters under oxygen to produce organo-mineral materials that have imbued properties which compare more favorably than the prevailing degraded forest soil nutrients. This opinion is in line with the articulations of [9] and [10] that combined application of organic matter and inorganic fertilizers is essential for optimum crop production and arrest of soil nutrient depletion in West Africa. Optimum plant media for germination and early seedling do not only require water but has the capacity to retain sufficient moisture, supply nutrients, and produce adequate aeration as any condition short of these leads to poor germination of seeds [11].

Although there is a general consensus that significantly more carbon store in the world soils exceeds that present in the atmosphere, [12] the availability of these soil organic matter for plant growth through decomposition is vital. Current debate on the sensitivity of organic matter decomposition to changes in temperature is therefore in order [13,14,15,16] particularly in view of declining forest vegetation and litter yield. It is against this backdrop that this study was conducted to evaluate the potency of topsoil from the readily available and fast growing exotic *Gmelina arborea* plantation by modification with temperature and organic matters as growth media for the early growth of *Irvingia gabonensis* for domestication purposes.

## 2. METHODS AND MATERIALS

### 2.1 Description of Study Area

This study was carried out in the Professor Nnabuife Screen House at the Department of Forestry and Wildlife, Faculty of Agriculture, Nnamdi Azikiwe University Awka, Anambra State. The University is located in the eastern part of Nigeria and lies between latitudes 6°06'N and 6°16'N, longitudes 7°01'E and 7°10'E.

Climatic condition of the area is tropical, dominated by rainfall pattern ranging from 1828 mm – 2002 mm. The average annual temperature and relative humidity is approximately 26.3°C and 80% respectively [17].

## 2.2 Media Procurement and Preparation

Litter-free top soil was technically collected from a litter-raked forest surface floor area of 5 x 5 m at a depth of 0-20 cm in the *Gmelina arborea* forest plantation within the Nnamdi Azikiwe University, Awka. The river sand was collected directly from a dredger in the River Niger at the Onitsha axis while the poultry floor waste was obtained from a local poultry in Awka.

The control, topsoil (T1) and river sand were then sieved with a 2 mm mesh size filter while the river-sand was further sterilized at 160°C for 24 hours in an oven. The topsoil was then primed by mixing with poultry floor waste and sterilized river sand in a uniform ratio of 4:3 to produce T2 and T3 soil amendments respectively.

The forest floor litters from the *Gmelina* plantation were screened for metals before it was sun-dried at screen house temperature 25-30°C for 48 hours and then combusted in the presence of minimal oxygen with underlying topsoil in relatively covered metallic buckets in the ratios 5:1, 10:1, 15:1 and 20:1 of litter to soil for 5, 10, 15 and 20 minutes to produce T4, T5, T6 and T7 thermally amended soils respectively.

The different soil amendments were sampled and analyzed for pH, Nitrogen, Calcium, Magnesium, Sodium, Potassium and Cation exchange capacity (CEC). Soil pH was determined in water by means of Bechman's pH meter using a soil to water ratio of 1:2.5 [18]. Exchangeable basic cations ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$  and  $\text{Na}^+$ ) were extracted with 1N  $\text{NH}_4\text{OAc}$  at pH 7; Ca and Mg were determined by Atomic Absorption Spectrophotometer (AAS) while K and Na were determined by Flame Photometry [19]. Cation exchange capacity was measured using ammonium acetate leaching at pH 7.0 [20]. Exchangeable acidity was extracted with normal KCl solution. The exchangeable acidity and exchangeable Aluminum were determined by titrate as described by [18]. The exchangeable hydrogen was obtained by subtracting exchangeable acidity (Al+H) – Exchangeable Al= Exchangeable H. Total nitrogen was determined by the micro Kjeldahl distillation method [21].

## 2.3 Fruit Preparation and Germination

Five hundred and fifty (550) ripe fruits of *Irvingia gabonensis* procured from the Department of Forestry of Anambra State Ministry of Environment were de-pulped manually with knife to scrap off the mesocarp. The obtained seeds were subjected to viability test by flotation method [22] and five hundred and forty three (543) that passed the test were sun dried at the screen house temperature of 25-30°C for 24 hours.

Seventy (70) seeds were then sown in seven open wooden germination beds measuring 2.5 m x 1.5 m x 0.3 m filled with the 6 different soil amendments and the control of unamended top soil. These were watered once with 5 liters of water by sprinkling and monitored for germination at 3 and 6 weeks after sowing respectively. Seed germination was obtained by observation of the viable protrusion on the soil surface of at least 0.50 cm of the cotyledon and hypocotyls of the seedlings.

At the end of 6 weeks, forty (40) sturdy seedlings each were pricked from promising amendments into uniform poly pots of 20 x 10 x 10 cm<sup>3</sup> containing the respective soil treatments and watered with 5 liters of water daily for 7 days under screen house conditions. These were thereafter monitored for early growth variables for a period of 6 weeks. The stem height was obtained with a steel meter rule from the soil level to the terminal bud while the leaf area was determined using portable leaf area meter (L1-COR 300°C). The collar diameter was measured with the digital vernier caliper while the number of leaf per plant was obtained by counting the number of leaves per pot per treatment.

Data collected were analyzed with the analysis of variance (ANOVA) and significant means separated at 5% level of significance with the Duncan multiple range test.

## 3. RESULTS

### 3.1 Characteristics of Soil Used in the Study

The chemical properties of soil amendments used in the study are as shown in Table 1. The highest Ca content was recorded by the T5 (40.67 meq/100 g soil) while the T7 (13.16 meq/100 g soil) was least. The control topsoil (T1) had calcium (Ca) content of 31.67 meq/100

g soil while the T2 (29 meq/100 g soil) and T3 (23.34 meq/100 g soil).

The magnesium content was highest in T5 (12.5 meq/100 g soil) while T7 (2.80 meq/100 g soil) recorded the least. The T4 (6 meq/100 g soil) and T1 (5 meq/100 g soil) recorded higher content than T2 (4.17 meq/100 g soil), T6 (3.83 meq/100 g soil) and T3 (3.33 meq/100 g soil). Sodium contents ranged from 0.11 meq/100 g soil in T4 to 0.28 meq/100 g soil in T3.

The cation exchange capacity (CEC) ranged from 27.80 meq/100 g soil for the T6 (15M-FTS) to 62.12 meq/100 g soil (10M-FTS). The control treatment (T1) recorded a CEC of 56.91 meq/100 g soil while the T2 and T3 recorded CEC of 43.87 meq/100 g soil and 31.29 meq/100 g soil respectively.

The exchangeable acidity (H<sup>+</sup> Al) was highest for T1 (19.75 meq/100 g soil) and least for T6 (1.75 meq/100 g soil). The EA for T2 and T3 was 10.5 meq/100 g soil and 10.38 meq/100 g soil respectively. The T2 and T3 recorded total nitrogen contents of 0.23% and 0.19%

respectively while T7 was least. The soil reaction was in the order T7 (8.1) > T6 (8.0) > T5 (7.0) > T3 (6.9) T4 (6.6) > T1 (6.2) > T2 (5.9).

### 3.2 Effect of Soil Amendments on Germination

The germination of sown Irvingia seeds on the various amended soil at 3 and 6 weeks are shown (Table 2). At 3 weeks after sowing, T4 recorded the highest number of germination percentage of 72.5% while the T2 recorded a germination percentage of 57.5%. The T1, T3 and T5 recorded 42.5 percent each while T6 supported the least germination percentage (28.57%). The T7 did not support the germination of any seed.

However at 6 weeks after sowing, the T4 maintained the highest germination percentage of 100% as all the 40 sown seeds germinated. The T1, T2 and T5 supported germination to the same capacity of 85.5% while the least germination percentage of 42.5% was shown by T6 at 6 weeks. The T7 recorded zero germination at 3 weeks and a germination percentage of 15% at 6 weeks.

**Table 1. Chemical characteristics of soil amendments**

Variables	Ca	Mg	Na	K	H <sup>+</sup>	Al <sup>3+</sup>	CEC	N (%)	pH (H <sub>2</sub> O)	Ca/Mg ratio
	meq/100 g soil									
T1	31.67	5.00	0.17	0.32	9.25	10.5	56.91	0.15	6.2	6.33
T2	29.00	4.17	0.18	0.27	4.50	5.75	43.87	0.23	5.9	6.95
T3	23.34	3.33	0.28	0.59	2.00	8.38	31.29	0.16	6.9	7.01
T4	25.83	6.00	0.11	0.36	6.25	8.25	46.81	0.19	6.6	4.31
T5	40.67	12.50	0.14	0.31	4.00	4.50	62.12	0.17	7.0	3.25
T6	22.24	3.83	0.14	0.34	1.50	0.25	27.80	0.14	8.0	5.81
T7	13.60	2.80	0.18	0.28	1.75	1.50	44.87	0.11	8.1	4.86

Key: T1= Topsoil; T2=Topsoil + poultry waste; T3= Topsoil + river-sand; T4= 5 mins combusted (topsoil + Gmelina forest floor litter); T5= 10 mins combusted (topsoil + Gmelina forest floor litter); T6= 15 mins combusted (topsoil + Gmelina forest floor litter); T7= 20 mins combusted (topsoil + Gmelina forest floor litters)

**Table 2. Effect of soil amendments on germination of Irvingia seeds**

Variables	No of germinated seeds		Germination percentage (%)	
	3 weeks	6 weeks	3 weeks	6 weeks
T1	17	34	42.5	85
T2	23	34	57.5	85
T3	17	29	42.5	72.5
T4	29	40	72.5	100
T5	17	34	42.5	85
T6	11	17	27.5	42.5
T7	0	1	0	15

Key: T1= Topsoil; T2=Topsoil + poultry waste; T3= Topsoil + river-sand; T4= 5 mins combusted (topsoil + Gmelina forest floor litter); T5= 10 mins combusted (topsoil + Gmelina forest floor litter); T6= 15 mins combusted (topsoil + Gmelina forest floor litter); T7= 20 mins combusted (topsoil + Gmelina forest floor litters)

**Table 3. Effects of soil amendments on Growth variables of *Irvingia gabonensis* seedlings**

Amendments	Stem height	Collar diameter	Leaf area	Leaf number
T1	97.1±0.79 <sup>b</sup>	3.20×10 <sup>-1</sup> ±0.03 <sup>c</sup>	20.66±2.85 <sup>d</sup>	3.06±0.33 <sup>a</sup>
T2	65.1±1.07 <sup>e</sup>	2.80×10 <sup>-1</sup> ±0.51 <sup>d</sup>	14.09±3.44 <sup>e</sup>	2.00±0.43 <sup>bc</sup>
T3	117.9±0.42 <sup>a</sup>	4.50×10 <sup>-1</sup> ±0.05 <sup>a</sup>	35.08±4.85 <sup>a</sup>	3.39±0.24 <sup>a</sup>
T4	89.8±0.71 <sup>c</sup>	3.80×10 <sup>-1</sup> ±0.05 <sup>b</sup>	22.75±4.09 <sup>c</sup>	2.78±0.25 <sup>b</sup>
T5	69.1±0.41 <sup>d</sup>	1.90×10 <sup>-1</sup> ±0.22 <sup>e</sup>	25.94±5.44 <sup>b</sup>	2.39±0.34 <sup>b</sup>
Mean	87.8±3.93	3.20×10 <sup>-1</sup> ±0.19	23.68±18.68	2.72±1.43
SE	0.41	0.02	1.98	0.15

Means ±Std. deviation with the same superscript and column are not significantly different ( $P < 0.05$ )

### 3.3 Effect of Amendments on Growth Variables

The results showed that T3 amendment recorded the highest height of 117.9±0.42 mm while the T2 least height (6.51±1.07 mm) as shown in Table 3. There were significant differences in all the effects of treatments on the height of *Irvingia*.

There was no significant difference in the leaf number of T3 and T1 as well as between T4, T5 and T2. The highest leaf number was recorded by T3 (3.39±0.24) while the least by T2 (2.00±0.43). There were significant differences in the leaf areas of seedlings and was in the order T3 (35.08±4.85 mm<sup>2</sup>) > T5 (25.94±5.44 mm<sup>2</sup>) > T4 (22.75 ±4.09 mm<sup>2</sup>). The least leaf area (14.09± 3.44 mm<sup>2</sup>) was recorded by T2.

There was also significant difference ( $P < 0.05$ ) in the collar diameters of seedlings as influenced by the various growth media. The order of influence was T3 (4.50×10<sup>-1</sup> ± 0.05 mm) > T4 (3.80×10<sup>-1</sup> ±0.05 mm) > T1 (3.20×10<sup>-1</sup> ± 0.03 mm) > T2 (2.80×10<sup>-1</sup> ± 0.51 mm). The least collar diameter of 1.90×10<sup>-1</sup> ± 0.22 mm was recorded by T5.

## 4. DISCUSSION

The germination potential of T4 and T5 which are temperature amended media producing the highest germination percentages at 3 and 6 weeks may not be unrelated to the resultant enhanced moisture content of the media. This is because of the ability of such media to hold and supply water to the seeds to imbibe, activate the enzymatic process needed to initiate seed germination [23] as well as promoting better root development to facilitate germination. This may also be due to a probable release of mineral elements as a result of the thermal treatment which may have solicited the release of mineral elements from combusting litters to form organo-mineral compounds that have the potential to break seed dormancy faster than sole organic

materials. The soil reaction values clearly support this assertion as the thermally modified amendments showed a two spectrum pH range from neutral (T4 and T5) to moderately alkaline (T6 and T7) that are quite different from the organically primed media.

The neutral pH range materials was able to attain 100% germination while T3 (topsoil + riversand) with strongly alkaline status gave 72.5% germination. This result did not only support [24] that recorded great success with sand +topsoil for seed germination of *Irvingia wombulu* and [25] that reported highest sprouting from top-river soil, but further advanced thermally modified media as a potentially better germination treatment media for the Irvingiaceae family.

The presence of organo-mineral elements accounted for significantly high cation exchange capacity (CEC) of T4 and T5 which compared favorably with T1 and T2 that recorded the same percentage germination percentage at 6 weeks after sowing. The CEC of thermally amended media increased with time from T4 to T5 and then fell at T6. This finding agrees with that of [26] which revealed that interactions between organic matters and soil particles increases the CEC with age and time to produce a range of functional groups at the soil surfaces.

The performance of T6 and T7 extended thermally treated soil media was abysmally poor. This may not be unexpected as according to [27,28] depending the timing and magnitude of thermal exposure of soils, depletion of microbial accessible carbon pools, microbial reduction as well as a shift in microbial use efficiency and modification of microbial community composition results which ultimately will result in poor seed germination rate and seedling performance. Subsequently, T6 and T7 were not considered promising enough to warrant trials in the growth variable experiment even though T6 recorded 27.5% and 42.5% at 3 and 6 weeks respectively. This poor performance may also not be unrelated

to the wide time lag which may have degraded the originally resultant organo-mineral elements attained at T4 and T5 time periods. In fact, the moderately alkaline (7.9-8.4) pH of T6 and T7 support this finding because alkaline conditions do not favor germination of tree crop species [29]. The T3, T4 and T5 were neutral (6.6-7.3) on the pH scale and recorded germination percentages of over 70%. The pH inclination of T3 towards thermally amended media which could equally have contributed to its efficacy may be as a result of the initial sterilization. But the biologically charred forest floor litters may have significantly accounted for the difference in performance.

The initial nitrogen content of T1 (0.15%) may have been enhanced by priming with poultry floor waste [30,31] and river-sand as well as the thermal modification of the media. However, the nitrogen content of thermally amended media reached maximum enhancement at T4 which later declined with increase heating time, becoming seemingly denitrified and immobilized at T6 and T7 with significant reduction on the exchangeable acidity (Al + H). This is because organic amendments can be used as a sink for reducing the bioavailability of metallic colloids in soils [32], which in proportionate term was highest in the 20minutes combusted Gmelina forest floor litter-soil (T7).

However, in the growth variable trials the T3 performed best in the stem height, collar diameter and leaf area. This may be due to its high potassium content which is important in plant metabolism, protein synthesis and chlorophyll development [33] and may also have accounted for the T4 and T5 that were next to T3 in the growth of collar diameter and leaf area respectively. Apart from the leaf numbers, the differences in the growth variables performances of T4 and T5 which differed significantly ( $P < 0.05$ ) can be attributed to the nutrient retention capacity that seem to be particularly higher in T4. This may not be unconnected with the wider Ca/Mg ratio of T4 in comparison with T5 indicating lower chances of leaching and reduction in the possibility of calcium deficiency for meristematic cell elongation and division. The Ca/Mg ratio of the organic amendments, T2 and T3, equally followed the same trend with wider ratio than the thermal amended soil media which perhaps conferred them with greater efficacies. The widest Ca/Mg ratio of T3 accounted for the highest performance of the growth variables.

## 5. CONCLUSION

The results of this study established that germination and early growth of *Irvingia gabonensis* depended on the Gmelina forest topsoil primed with river sand and the 5minutes thermally amended Gmelina plantation floor litter. It showed that the thermally amended soils led to the early and 100% germination of seeds while the river sand-primed amendment had greater influence on the growth variables although the former media compared favorably. These two types of soil media performed better than the traditional control top soil because of the capacities to hold water for protoplasmic activities and retain nutrients for vegetative development respectively, thereby greatly minimized leaching. Therefore, the domestication of this near threatened and economic forest tree species is possible with a combination of these cheap and readily available amendments at the nursery and plantation establishment phases.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Dolor DE. Propagation of *Treculia africana* as influenced by seed storage and propagation media. *Agricultura Tropica et Sub-tropica*. 2013;46(2):52-57.
2. Adelani DO, Suleiman RA, Olaiya RK, Yohanna EA. Hormonal pretreatments of African locust bean tree seeds (*Parkia biglobosa*, Jacqubenth). In: *Sudano-sahelian Landscape and Renewable Natural Resources Development in Nigeria*. Ogunsanwo, O.Y., Akinwale, A.O., Azeez, I.O., Adekunle, V.A.J. and Adewole, N.A. (eds); Proceedings of the 37<sup>th</sup> Annual Conference of Forestry Association of Nigeria. 2014;337-346.
3. Egwunatum AE. Sustainable use of forest wealth: Critical challenges, controversies and conservation. In: *Forest: Its wealth and Future*. Akpan, M. and Udofia, S.I. (eds); Proceedings of the National Workshop of The Forestry Association of Nigeria, Akwalbom State Branch. 2018;129-134.
4. Bargali K, Tewari A. Growth and water relation parameters in drought-stressed *Coriaria nepalensis* seedlings. *J. Arid Environ*. 2004;58:505-512.

5. Burkil HM. The useful plants of West Tropical Africa. 2<sup>nd</sup> edition. Royal Botanic Gardens, Kew, UK. 2004;960.
6. Nya PJ, Omokaro DN, Nkang AE. Comparative studies on seed morphology, moisture content and seed germination of two varieties of *Irvingia gabonensis*. Global Journal of Pure and Applied Science. 2000;12(2):141-148.
7. Okafor JC. Checklist of medical plants of Nigeria and their uses. Association of Scientific Identification, Conservation and Utilization of Medicinal Plants of Nigeria (Asicumon) Enugu, Nigeria. Janoe Publishers; 2005.
8. IUCN. Red list of *Threatened* species. International Union for Conservation of Natural Resources: The Gymnosperm database- Conservation Vulnerable Status; 2006.
9. FAO. Soil fertility initiative for Sub-Saharan Africa. Proc. SFI/FAO consultation, Rome.19-20 Nov. 1999. FAO, Rome; 1999.
10. Giller KF. Targeting management of organic resources and mineral fertilizers: Can we match Scientists fantasies with Farmers realities? In: Vanlauwe, B., Diels, J., Saginga, N. and Merckx, R. (eds) Balanced Nutrient Management Systems for the Moist Savanna and Humid Forest Zones of Africa. CAB International, Wallingford, UK. 2002;155-171.
11. Ahmad S, Ahmad R, Ashraf MY, Ashraf M, Waraich EA. Sunflower (*Helianthus annuus*, L.) response to drought stress at germination and seedling growth stages. Park J. Bot. 2009;41(2): 647-654.
12. Sulman BN, Moore JAM, Abramoff R, Averill C, Kivlin S, Georgiou K, Sridhar B, et al. Multiple models and experiments underscore large uncertainty in soil carbon dynamics. Biogeochemistry. 2018;141(2):109–123.
13. Davidson E, Janssens I. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. Nature. 2006;440:165–173.
14. Xu Wenfang, Wenping Yuan, Lilun Cui, Minna Ma, Fenguo Zhang. Responses of soil organic carbon decomposition to warming depend on the natural warming gradient. Geoderma. 2019;343:10-18.
15. Neal E. Flanagan, Hongjun Wang, Scott Winton, Curtis J. Richardson. Low-severity fire as a mechanism of organic matter protection in global peatlands: Thermal alteration slows decomposition. Global Change Biology. 2020;1–17.
16. Wang JY, Ren CJ, Feng XX, Zhang L, Doughty R, Zhao FZ. Temperature sensitivity of soil carbon decomposition due to shifts in soil extracellular enzymes after afforestation, Geoderma. Geoderma. 2020;374(2020):114426.
17. [NiMet] Nigerian Meteorological Agency. Quarterly weather bulletin; 2016. [Accessed 2017 May 29] Available:<https://nimet.gov.ng/sites/default/files/publications/2016%20Quarterly%20Bulletin.pdf>
18. Thomas GW. Exchange cations. IN. Page AI, Miller PH, Keeney DR. (Eds), Methods of soil analysis Part 2 Agro. Monogr. 9. 2<sup>nd</sup> ed Madison. 1982;159-165.
19. Jackson ML. Soil chemical analysis. Prentice Hall, Englewood Cliffs; 1958.
20. Rhoades JD. Cation exchange capacity in Page AL, Miller RH, Keeney DR. (eds). Methods of Soil Analysis Part 2. American Society of Agro, Madison, USA. 1982;149-158.
21. Bremner JM. Nitrogen-total. In: Chemical Methods of Soil Analysis Part 3 (eds). Sparks, D.L., Page, A.L., Helmke, P.A., Loeppert, R.H., Soltanpour, P.N., Tabatabai, M.A., Johnson, C.T. and Summer, M.E.). Soil Science Society of America Book series, American Society of Agronomy, Madison USA. 1996;1085-1121.
22. Daneshvar A, Tigabu M, Karimidoost A, et al. Flotation techniques to improve viability of *Juniperus polycarpus* seed lots. J. For. Res. 2017;28:231–239.
23. Ekebafé MO, Oviasogie PO, Ekebafé LO. Effect of palm fronds and cow dung biochar on the soil-water retention capacity and growth of oil palm (*Elaeis guineensis*) sprouted seedlet. In: Soil resources management, Global climate change and Food security. Asadu CLA, Ezeaku PI, Uzoh MI, Unagwu B (eds). Proceedings of the 36<sup>th</sup> Annual Conference of Soil Science Society of Nigeria. 2012;127-139.
24. Dolor D. Effect of propagation media on the germination and seedling performance of *Irvin giawombulu* (Vermoesen). American Journal of Biotechnology and Molecular Sciences. 2011;1(2):51-56
25. David J, Shimber T, Amsalu N. Effect of nursery potting media and watering frequency on emergence and seedling growth of Korarima (*Aframomum corrorima*

- (Braun), Sky Journal of Agricultural Research. 2014;3(10):187-195.
26. Cheng Chi-Hsin, Johannes Lehmann, Janice E. Thies, Sarah D. Burton, MMark H. Engelhard. Oxidation of black carbon by biotic and abiotic processes. Organic Geochemistry. 2006;37(11): 1477-1488.
27. Melillo JM, Butler SM, Johnson JE, Mohan J, Steudler PA, Lux H, Burrows E, Bowles F, Smith RM, Scott L, Vario CL, Hill TD, Burton AJ, Zhou Y, Tang J. Soil warming, carbon-nitrogen interactions and forest carbon budgets. Proc. Natl. Acad. Sci. 2010;108:9508–9512.
28. Melillo JM, Frey SD, DeAngelis KM, Werner WJ, Bernard MJ, Bowles FP, Pold G, Knorr MA, Grandy AS. Long-term pattern and magnitude of soil carbon feedback to the climate system in a warming world. Science. 2017;358(6359):101–105.
29. Acquash G. Principles of crop production: Theory, techniques and technology. Prentice-Hall of India Private Limited, New Delhi. 2005;165-205.
30. Acosta-Martinez V, Harmel RD. Soil microbial communities and enzyme activities under various poultry litter application rates. Journal of Environmental Quality. 2006;35:1309-1318.
31. Adeli A, Tewolde H, Sistani KR, Rowe DE. Broiler litter fertilization and cropping system impacts on soil properties. Agronomy Journal. 2009;110:1304-1310.
32. Park JH, Lamb D, Paneerselvam P. Role of organic amendments on enhanced bioremediation of heavy metalloids contaminated soils. Journal of Harzardous Materials. 2011;185:549-574.
33. Nwoboshi LC. The nutrient factor in sustainable forestry. Ibadan University Press, Nigeria. 2000;303.

© 2020 Egwinatum et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:  
<http://www.sdiarticle4.com/review-history/58259>*